
Stereoselective synthesis of mexiletine and structural analogs with chiral tert-butanefulfinamide

Journal: Tetrahedron Letters

Publication Year: 2015

Authors: Daniel A. Ryan, Karl J. Ololotowicz, Mark Mercola, John R. Cashman

PubMed link:

Funding Grants: Improving Existing Drugs for Long QT Syndrome type 3 (LQT3) by hiPSC Disease-in-Dish Model

Public Summary:

An asymmetric synthesis of mexiletine and structural analogs was developed using chiral tert-butanefulfinamide to convert precursor ketones to chiral amines. Starting from α -aryloxy ketones, a two-step condensation–reduction procedure provided chiral N-tert-butanefulfinyl amines as immediate precursors to mexiletine or structural analogs. Reduction of the intermediate N-tert-butanefulfinyl imine showed substrate- and reagent-derived stereoselectivity. Following removal of the chiral auxiliary, mexiletine and structural analogs were obtained in high enantiopurity using this approach.

Scientific Abstract:

An asymmetric synthesis of mexiletine and structural analogs was developed using chiral tert-butanefulfinamide to convert precursor ketones to chiral amines. Starting from α -aryloxy ketones, a two-step condensation–reduction procedure provided chiral N-tert-butanefulfinyl amines as immediate precursors to mexiletine or structural analogs. Reduction of the intermediate N-tert-butanefulfinyl imine showed substrate- and reagent-derived stereoselectivity. Following removal of the chiral auxiliary, mexiletine and structural analogs were obtained in high enantiopurity using this approach.

Source URL: <https://www.cirm.ca.gov/about-cirm/publications/stereoselective-synthesis-mexiletine-and-structural-analogs-chiral-tert>